



UNIVERSITI PUTRA MALAYSIA

**THE PRESENCE OF STAPHYLOCOCCUS AUREUS ON THE SKIN,
NOSE, AND EAR OF ADOLESCENT FOOTBALLERS BEFORE AND
AFTER TRAINING**

JOHN LAURENCE WILLIAM.

FPP 2005 17

THE PRESENCE OF *STAPHYLOCOCCUS AUREUS* ON THE SKIN,
NOSE, AND EAR OF ADOLESCENT FOOTBALLERS BEFORE AND
AFTER TRAINING

By

JOHN LAURENCE WILLIAM

Thesis Submitted to the School of Graduate Studies, University Putra
Malaysia, in Fulfilment of the Requirements for the Degree of Doctor of
Philosophy

October 2004

DEDICATION

**To My Late Father, Beloved Mother, Family Members, My Dearest Wife
And Children Jasmeen, Josephine, Jemima, Jarvis**

Abstract of thesis presented to Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Doctor of Philosophy

THE PRESENCE OF *STAPHYLOCOCCUS AUREUS* ON THE SKIN,
NOSE, AND EAR OF ADOLESCENT FOOTBALLERS BEFORE AND
AFTER TRAINING

By

JOHN LAURENCE WILLIAM

October, 2004

Chairman : Saidon Amri, PhD

Faculty : Educational Studies

The objective of the study was to determine the number of adolescent footballers carrying *Staphylococcus aureus* during outdoor and indoor training. One hundred twenty healthy male subjects age from 13 to 15 participated in the study. The subjects were assigned into two different training venues; outdoors and indoors. Swabs were taken before and after training at three sites of the body; skin, ear and nose. The strains were incubated in Baird Parker agar plates at 37°C. There was no significant difference for the numbers of *S. aureus* strains carriers between pre and post result for the skin, nose, and ear for outdoor training.

Results for the number of *S. aureus* strains carriers on the skin, and nose between pre and post indoor training showed significant difference ($p<0.05$), however no significant difference was observed for the results on the ear. Number of carriers on the skin, ear, and nose for pre training was 60 (100%), 56 (93%) and 56 (93%), respectively, whereas for post training on the same sites were 44 (73%), 56 (93%), and 60 (100%), respectively. As for the estimated mean plate count of *S. aureus* strains, there was significant difference between the pre and post results on the ear and nose of Malays during outdoor training ($p<0.05$) and no significant difference for the estimated mean plate count of strain on the skin. The estimated mean plate count of *S. aureus* strain for Malays from the skin, ear and nose before training were 138 ± 103 , 190 ± 147 and 395 ± 83 , respectively and after training were 97 ± 77 , 71 ± 64 and 498 ± 75 , respectively. For Indians, there was no significant difference for the plate count of *S. aureus* strain on the skin, ear, and nose between the pre and post results during outdoor training. As for indoor training, both Malays and Indians showed a significant difference between the pre and post results ($p<0.05$). The estimated mean plate count of strain during indoor training for Malays from the skin, ear and nose before training was 33 ± 17 , 71 ± 28 and 312 ± 55 , respectively and 21 ± 16 , 44 ± 26 and 452 ± 89 , respectively after training. For Indians, the estimated mean plate count of strain on the skin, ear, and nose before training was 72 ± 21 , 80 ± 21 and

309±104 respectively and 55±19, 200±62, and 466±109 respectively after training. In conclusion, the adolescent footballers are carriers during training either indoor or outdoor.

Representative strains from the skin of adolescent footballers and environment were selected randomly for antibiotic resistance, plasmid, coagulase, and RAPD-PCR analysis. For the antibiotic resistance test, nineteen antibiotics were tested. Antibiotic resistance patterns with the strains tested from footballers training indoor, outdoor and environment were diverse. However, norfloxacin (0%), rifampicin (0%), imipenem (0%), methicillin (0%) and trimethoprim-sulfamethoxazole (100%) showed a similar resistance patterns with the strains tested from footballers and the environment. Strains from the adolescent footballers training indoors and indoor environment have a higher Multiple Antibiotic Resistance (MAR) index compared from the strains from outdoor adolescent footballers and outdoor environment. The plasmid profiles of *S. aureus* strains isolated from the skin of footballers and the environment ranged between 1.8 to 3.4 megaDalton (mDa). The results of the plasmid profiles and antibiotic resistance showed that there was no correlation between plasmid carriage and resistance to a particular antibiotic tested. Thirty-two *S. aureus* strains isolated from outdoors and indoors were found to be carrying coagulase genes of different sizes. Two strains produced three

amplified coagulase gene fragments while one strain produced two amplified coagulase gene fragments. The rest of the strains produced only one amplified coagulase gene fragment. As for the strains from the environment, two strains produced two amplified coagulase gene fragments and seven produced only one amplified coagulase gene fragment. The profiles obtained from RAPD contained 1 to 9 bands within the molecular size of 0.3 kbp to 5.0 kbp. From the dendrogram, the strains were divided into 2 major clusters and at 100% similarity there were four groups of strains. It can be concluded that the strains isolated from the footballers and environment were pathogenic due to the present of the coagulase gene.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

KEHADIRAN *STAPHYLOCOCCUS AUREUS* PADA KULIT, HIDUNG,
DAN TELINGA PEMAIN REMAJA BOLASEPAK SEBELUM DAN
SELEPAS LATIHAN

Oleh

JOHN LAURENCE WILLIAM

Oktober, 2004

Pengerusi : Saidon Amri, PhD

Fakulti : Pengajian Pendidikan

Objektif kajian adalah untuk menentukan jumlah pemain remaja bolasepak pembawa *Staphylococcus aureus* semasa menjalani latihan di dalam dewan dan padang. Seratus dua puluh subjek berumur di antara 13 hingga 15 tahun terlibat dalam kajian ini. Subjek-subjek di bahagikan kepada dua tempat latihan yang berbeza; di luar padang dan dalam dewan. Swabs diambil sebelum dan selepas latihan pada tiga bahagian badan; kulit, telinga, and hidung. Kesemua swabs di kulture dengan agar Baird Parker dan dieram pada suhu 37°C.

Semasa latihan di padang tidak ada kesan signifikan pada jumlah pembawa *S. aureus* sebelum dan selepas latihan pada kulit, hidung and telinga. Sebaliknya semasa latihan di dewan terdapat kesan signifikan ($p<0.05$) pada jumlah pembawa *S. aureus* sebelum dan selepas latihan pada kulit dan hidung, manakala tidak ada kesan signifikan pada jumlah pembawa pada telinga. Purata pembawa *S. auerus* pada kulit, hidung, and telinga sebelum latihan adalah 60 (100%), 56 (93%) dan 56 (93%), manakala selepas latihan ialah 44 (73%), 56 (93%) dan 60 (100%) masing-masing. Terdapat kesan signifikan jumlah anggaran *S. aureus* pada telinga dan hidung pemain Melayu semasa latihan di padang dan tiada kesan signifikan jumlah *S. aureus* pada kulit ($p<0.05$). Jumlah anggaran *S. aureus* pada kulit, telinga dan hidung adalah 138 ± 103 , 190 ± 147 dan 395 ± 83 sebelum latihan, dan selepas latihan adalah 97 ± 77 , 71 ± 64 and 498 ± 75 . Untuk pemain India tiada kesan signifikan jumlah anggaran *S. aureus* semasa latihan di padang pada kulit, hidung dan telinga. Untuk latihan di dewan, terdapat kesan signifikan jumlah anggaran *S. aureus* untuk Melayu dan India pada kulit, telinga, dan hidung ($p<0.05$), Jumlah anggaran *S. aureus* sebelum latihan untuk pemain Melayu adalah 33 ± 17 , 71 ± 28 dan 312 ± 55 , manakala selepas latihan adalah 21 ± 16 , 44 ± 26 , 452 ± 89 masing-masing. Untuk pemain India jumlah anggaran *S. aureus* pada kulit, telinga, dan hidung semasa latihan

dalam untuk pra dan post latihan adalah 72 ± 21 , 80 ± 21 , 309 ± 104 , dan 55 ± 19 , 200 ± 62 , 466 ± 109 masing-masing.

Strain yang mewakili dari kulit subjek dan udara di pilih secara rambang untuk ujian kerintangan antibiotik, plasmid, coagulase and RAPD-PCR. Strain dari kulit subjek dan udara diuji kerintangan antibiotik dengan sembilanbelas antibiotik. Keputusan kerintangan antibiotik untuk strain dari subjek luar, dalam dan udara mempunyai corak yang berbeza. Walaupbagaimanapun antibiotik norfloxacin (0%), rifampicin (0%), imipenem (0%), methicillin (0%) and trimethoprim-sulfamethoxazole (100%) menunjukkan corak kerintangan yang sama bila diuji dengan strain dari subjek dan udara. Nilai MAR index adalah tinggi untuk strain dari subjek dalam dan udara dalam. Secara kesimpulan strain ini adalah lebih patogenik jika dibandingkan dengan strain lain.

Keputusan menunjukkan bahawa strain dari kulit subjek dan udara mengandungi plasmid dengan saiz di antara 1.8 hingga 3.4mDa. Bagaimanapun keputusan tidak menunjukkan sebarang perkaitan di antara profil plasmid dengan profil ketahanan antibiotik tertentu. Tiga puluh dua strains dari kulit subjek semasa latihan di luar dan dalam di dapati membawa gen koagulase dengan saiz yang berbeza. Dua strain dari subjek di dapati mempunyai tiga fragmen gen koagulase and satu

strain mempunyai dua fragmen gen koagulase. Strain yang lain hanya mempunyai satu fragmen gen koagulase. Dua strain dari udara mempunyai dua fragment gen koagulase dan yang lain satu fragment gen koagulase. Keputusan profil 'fingerprinting' yang diperolehi dari RAPD mengandungi 1 hingga 9 jalur dengan saiz molekul 0.3 kbp hingga 5.0 kbp yang membentuk dua congkok utama. Pada 100% kesamaan terdapat empat congkok strain. Kesimpulannya strains daripada subjek and udara adalah patogenik kerana membawa gen koagulase.

ACKNOWLEDGEMENTS

Firstly I would like to take this opportunity to thank all the people concerned for helping and guiding me during my study. With their help and support, I finally managed to complete my study.

I would like to express my sincere thanks and appreciation to my co-supervisor Professor Dr. Son Radu and Associate Professor Dr. Abdul Raha Abdul Rahim. Professor Son Radu who has devoted a lot of his time for me during my study. His guidance, advice, supervision and support have gave me a lot of knowledge and confidence. The knowledge and research, which he shares with me, has opened a new dimension in Sport Science. Associate Professor Raha Abdul Rahim too played an important role during my course. Her advice, and support have encouraged me towards the completion of this work and being helpful whenever I face problems. Also thanks to Dr. Saidon Amri for sharing some of his knowledge with me.

Sincere thanks also to Associate Professor Dr. Shahrauddin Abdul Aziz for his advice and helpful discussion that enlightened me to work harder and complete my study. His short period with me was really worth while and valuable. A special thanks to the School of Graduate Studies,

Professor Dr. Aini Ideris and Professor Dr. Gulam Rusul Rahmat Ali for their kind help to settle problems.

I would also like to thank the Faculty of Food Science and Biotechnology for allowing me to use the facilities and laboratory for my work. With the laboratory facilities I managed to finish my laboratory work.

Also, I had been very fortunate to have a very kind and understanding family members. My mother's advice always enlightens me whenever I feel I am moving towards the wrong direction. Thanks to my wife Munica and my children Jasmeen, Josephine, Jemima and Jarvis for their patience with me. It is really great to have all of you.

Lastly I would also like to thank to my friends Kqueen, Samuel, Audrey, Lesley, Rozsilah, and Ooi Wai Lieng for their knowledge and time that they shared with me. The knowledge I gained was valuable for my future research. For those who have helped me and whose names have not been mentioned, I am very sorry. It would not be possible to name all of them here but I will remember all your assistance during my study. May God Bless all of you.

I certify that an Examination Committee met on 29th October 2004 to conduct the final examination of John Laurence William on his Doctor of Philosophy thesis entitled "The Presence of *Staphylococcus aureus* on the Skin, Nose, and Ear of Adolescent Footballers Before and After Training" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Shamsher Mohamad Ramadili, PhD

Professor
Faculty of Economics and Management
Universiti Putra Malaysia
(Chairman)

Norhani Abdullah, PhD

Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Mariana Nor Shamsudin, PhD


Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Internal Examiner)

Michael Gleeson, PhD

Professor
Loughborough University
United Kingdom
(External Examiner)

N.C. Craig Sharp, PhD

Professor
Brunel University
United Kingdom
(External Examiner)



ZAKARIAH ABDUL RASHID, PhD
Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 JUL 2005


This thesis submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows :

Saidon Amri, PhD
Faculty of Graduate Studies
Universiti Putra Malaysia
(Chairman)

Son Radu, PhD
Professor
Faculty of Graduate Studies
Universiti Putra Malaysia
(Member)

Raha Abdul Rahim, PhD
Associate Professor
Faculty of Graduate Studies
Universiti Putra Malaysia
(Member)

Shahrauddin Abdul Aziz, PhD
Associate Professor
Faculty of Graduate Studies
Universiti Putra Malaysia
(Member)



AINI IDERIS, Ph.D.
Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia

Date : 11 AUG 2005

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



JOHN LAURENCE WILLIAM

Date : 15/7/2005

LIST OF TABLES

Table	Page
1.1 Infection of skin and soft tissue	10
2.1 Classification of physical activity intensity and RPE	85
4.1 Physical characteristics of the Malay and Indian subjects during outdoor and indoor training (n=120)	126
4.2 The number of adolescent footballers carrying <i>S. aureus</i> during outdoor and indoor training (n=120)	128
4.3 Estimated plate count of <i>S. aureus</i> during pre and post training for outdoor venue (n=60)	131
4.4 Estimated plate count of <i>S. aureus</i> during pre and post training at indoor venue (n=60)	133
4.5 Estimated plate number <i>S. aureus</i> and temperature from outdoor and indoor venue	134
4.6 Antibiotics agents testing for susceptibility and resistance with strains isolated from adolescent footballers during outdoor training.	137
4.7 Antibiotics agents testing for susceptibility and resistance with strains isolated from adolescent footballers during indoor training.	140
4.8 Antibiotics agents testing for susceptibility and resistance with strains isolated from environment outdoor and indoor training.	144
4.9 Antibiotic resistance patterns of <i>S. aureus</i> strains isolated from adolescent footballers during outdoor training	152
4.10 Antibiotic resistance patterns of <i>S. aureus</i> strains isolated from adolescent footballers during indoor training	153
4.11 Antibiotic resistance patterns of <i>S. aureus</i> strains isolated from outdoor and indoor environment	155



4.12	Antibiotic resistance patterns and plasmid profiles of <i>S. aureus</i> strain isolated from footballers during outdoor training.	157
4.13	Antibiotic resistance patterns and plasmid profiles among <i>S. aureus</i> strains isolated from footballers during indoor training.	160
4.14	Antibiotic resistance patterns and plasmid profiles of <i>S. aureus</i> strains isolated from environment.	162
4.15	Antibiotic resistance patterns and coagulase gene strains isolated from adolescent footballers during outdoor and indoor training.	170
4.16	Antibiotic resistance pattern and coagulase gene strains isolated outdoor and indoor environment.	171
4.17	<i>Staphylococcus aureus</i> strains isolated from adolescent footballers during outdoor training	172
4.18	<i>Staphylococcus aureus</i> strains isolated from adolescent footballers during indoor training	173
4.19	<i>Staphylococcus aureus</i> strains isolated from the environment	174

LIST OF FIGURES

Figure		Page
2.1	Grapelike clusters of <i>Staphylococcus aureus</i>	34
2.2	Sites of infection and diseases caused by <i>S. aureus</i>	35
2.3	Virulence factors of <i>S. aureus</i> that are toxin to the host	36
2.4	Coagulase triggers blood plasma clotting and allowing protection from the immune system	38
2.5	Damaged of heart valve cause by <i>S. aureus</i>	46
2.6	Meningitis is disease inflammation of the meninges cause by <i>S. aureus</i>	47
2.7	The activities of the chemical mediators of inflammation	58
2.8	The process of maturation for lymphocytes into B and T cell	60
2.9	The function of antibodies depending on the class of the antibody and the nature of the antigen	62
2.10	The 'Inverted J Hypothesis'	69
2.11	The five modes of action by which drugs exert their antimicrobial effects	74
2.12	Polymerase chain reaction (PCR)	88
3.1	Flow chart of the experimental design	97
3.2	Flow chart design of the tests	98
3.3	<i>S. aureus</i> black colour colonies on the Baird Parker agar	102
3.4	Flow chart for identification test	109
3.5	Antibiotics testing by the disk diffusion method	112

3.6	Flow chart of DNA extraction	117
4.2	Mean value of estimated plate count of <i>S. aureus</i> during pre and post for the Malay and Indian adolescent footballers during outdoor training	131
4.3	Mean value of estimated plate count of <i>S. aureus</i> during pre and post for the Malay and Indian adolescent footballers during indoor training (n=60)	133
4.4	Resistance percentage of <i>S. aureus</i> strains isolated from adolescent footballers during outdoor training	138
4.5	Resistance percentage of <i>S. aureus</i> strains isolated from adolescent footballers during indoor training	141
4.6	Comparison antibiotic resistance pattern of strains isolated from adolescent footballers during outdoor and indoor training.	142
4.7	Resistance percentage of <i>S. aureus</i> strains isolated from environment outdoor and indoor training	145
4.8	Comparison antibiotic resistance pattern of strains isolated from environment outdoor and indoor	146
4.9	Comparison of antibiotic resistance pattern of strains isolated from Malay and Indian adolescent footballers during outdoor training	148
4.10	Comparison of antibiotic resistance pattern of strains isolated from Malay and Indian adolescent footballers during indoor training	150
4.11	Agarose gel (0.8%) electrophoresis of plasmid DNA from <i>S. aureus</i> strains isolated from footballers during outdoor training. Lane 1 marker (M) and follow by strains correspond with numbers 1 to 12.	158
4.12	Agarose gel (0.8%) electrophoresis of plasmid DNA from <i>S. aureus</i> strains isolated from footballers during outdoor training. Lane 1 marker (M) and follow by strain correspond with numbers 13 to 23.	158

4.13	Agarose gel (0.8%) electrophoresis of plasmid DNA from <i>S. aureus</i> strains isolated from footballers during indoor training. Lane 1 marker (M) and follow by strains correspond with numbers 1 to 13	161
4.14	Agarose gel (0.8%) electrophoresis of plasmid DNA from <i>S. aureus</i> strains isolated from footballers during indoor training. Lane 1 marker (M) and follow by strains correspond with numbers 14 to 29	161
4.15	Agarose gel (0.8%) electrophoresis of plasmid DNA from <i>S. aureus</i> strains isolated from thr environment. Lane 1 marker (M) and follow by strains correspond with numbers 1 to 10	163
4.16	Coagulase gene profiles of <i>S. aureus</i> isolates from footballers during outdoor training with COAG2 and COAG3 primers electrophoresed on 1% agarose gel. M: DNA ladder mix marker and correspond with the isolates numbers. Lane 17 is the negative control	165
4.17	Coagulase gene profiles of <i>S. aureus</i> isolates from footballers during indoor training with COAG2 and COAG3 primers electrophoresed on 1% agarose gel. M: DNA ladder mix marker and correspond with the isolates numbers. Lane 17 is the negative control	165
4.18	Coagulase gene profiles of <i>S. aureus</i> isolates from the environment with COAG2 and COAG3 primers electrophoresed on 1% agarose gel. M: DNA ladder mix marker and correspond with the isolates numbers. Lane 17 is the negative control	166
4.19	RAPD profiles of <i>S. aureus</i> obtained with using primer GEN1-50-01(5'GTGCAATGAG-3') electrophoresed on 1% agrose gel. M: DNA ladder mix marker and correspond with the isolates numbers	168
4.20	RAPD profiles of <i>S. aureus</i> obtained with using primer GEN1-50-01(5'GTGCAATGAG-3') electrophoresed on 1% agrose gel. M: DNA ladder mix marker and correspond with the isolates numbers	168

4.21	Dendrogram generated from the RAPD-PCR analysis using primer GEN1-50-01 (5'GTGCAATGAG-3') of the <i>Staphylococcus aureus</i> strains.	169
------	--	-----

LIST OF ABBREVIATIONS

µl	Microliter
µg	Microgram
µm	Micrometer
Bp	Base pair
DNA	Deoxyribonucleic acid
dNTP	Deoxynucleotide triphosphate
g	Gram
HCl	Hydrochloric acid
KAc	Potassium acetate
mg	Milligram
MgCl ₂	Magnesium chloride
ml	Milliliter
mM	MilliMolar
mm	Millimeter
mol	Mole
NaCl	Sodium chloride
NaOH	Sodium hydroxide
°C	Degree celsius
PCI	Phenol-Chloroform-Isoamyl alcohol
PCR	Polymerase Chain Reaction
RAPD	Random Amplified Polymorphic DNA

RNA	Ribonucleic acid
Rpm	Revolution per minute
SDS	Sodium dodecylsulphate
<i>Taq</i>	<i>Thermus aquaticus</i> DNA
TBE	Tris-Borate EDTA
Tris	Tris methylamine
TSB	Tryptic Soy Broth
UV	Ultraviolet
V	Volts
W/v	Weight / Volume
VO _{2max}	Maximum oxygen uptake
RBC	Red blood cell
WBC	White blood cell
NK	Natural killer cell
HDL	High density lipoproteins
LDL	Low density lipoproteins
ATP	Adenosine triphosphate
BMR	Basal metabolic rate
URTI	Upper respiratory tract infection
RPE	Rating of perceived exertion

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vii
ACKNOWLEDGEMENTS	xi
APPROVAL	xiii
DECLARATION	xv
LIST OF TABLES	xvi
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xxi

CHAPTER

1 INTRODUCTION

1.1	Background Of The Study	1
1.2	Statement Of Problem	11
1.3	Objective Of The Study	13
1.4	Purpose Of The Study	14
1.5	Research Questions	14
1.6	Significance Of The Study	15
1.7	Scope And Limitations Of The Study	16
1.8	Operational Definitions	17

2 LITERATURE REVIEW

2.1	Exercise And Health	23
2.2	<i>Staphylococcus aureus</i>	30
2.3	<i>Staphylococcus aureus</i> Virulence Factors	35
2.4	<i>Staphylococcus aureus</i> Diseases	43
2.5	Blood And Cellular Elements	48
2.6	Body's Defence System	52
2.7	Specific Body Defence	59
2.8	Exercise And Immune System	63
2.9	Antibiotic	70
2.10	Antibiotic Resistance	75
2.11	Plasmid	79
2.12	Rating Perceived Exertion (RPE)	82